



TITLE:

Finite Type System of Partial Differential Operators and
Decomposition of Solutions of Partial Differential Equations
(位相解析的方法による偏微分方程式論研究会及び散乱理
論の数学研究会報告集)

AUTHOR(S):

MATSUURA, SHIGETAKE

CITATION:

MATSUURA, SHIGETAKE. Finite Type System of Partial Differential Operators and Decomposition of Solutions of Partial Differential Equations (位相解析的方法による偏微分方程式論研究会及び散乱理論の数学研究会報告集). 数理解析研究所講究録 1967, 22: 10-17

ISSUE DATE:

1967-05

URL:

<http://hdl.handle.net/2433/107478>

RIGHT:

Finite Type System of Partial
Differential Operators and
Decomposition of Solutions of
Partial Differential Equations^(*)

by

Shigetake MATSUURA

§1. Introduction, $\mathcal{P}(D)$ -convexity.

First we fix some notations. Let Ω be an open set in the n -dimensional Euclidean space \mathbb{R}^n whose points shall be denoted by their coordinates $x = (x_1, \dots, x_n)$. Let $\mathbb{C}[X_1, \dots, X_n]$ be the polynomial ring in n variables $X = (X_1, \dots, X_n)$ over the complex number field \mathbb{C} .

A partial differential operator with constant coefficients $P(D)$ is obtained from the polynomial $P(X) \in \mathbb{C}[X_1, \dots, X_n]$ just by replacing the variables $X = (X_1, \dots, X_n)$ by the differentiations $D = (D_1, \dots, D_n)$ with $D_j = \frac{\partial}{\partial x_j}$ ($j = 1, 2, \dots, n$).

Now let $\mathcal{P}(X) = (P_{jk}(X))$ be a matrix with q rows and p columns with coefficients in $\mathbb{C}[X_1, \dots, X_n]$. $\mathcal{Q}(X) = (Q_{jk}(X))$ be a relation matrix with r rows and q columns for $\mathcal{P}(X)$, i.e. the row vectors $(Q_{j1}(X), \dots, Q_{jq}(X))$ ($j = 1, \dots, r$) generate the relation module

(*) The content of this article was partly spoken in a slightly different form in Séminaire MALGRANGE (1964) à Orsay.

$$\{(Q_1(X), \dots, Q_q(X)) ; \sum_{j=1}^q Q_j(X) P_{jk}(X) = 0 \quad (k = 1, \dots, p)\} .$$

Then we get the following differential complex

$$(1.1) \quad [C^\infty(\Omega)]^p \xrightarrow{P(D)} [C^\infty(\Omega)]^q \xrightarrow{Q(D)} [C^\infty(\Omega)]^r ,$$

$$Q(D)P(D) = 0 .$$

DEFINITION. An open set Ω in \mathbb{R}^n is called $P(D)$ -convex if the above sequence is exact.

EXAMPLES.

1) For the case of a single operator $P(D)$ ($p = q = 1$) , the exactness of (1.1) is reduced to the subjectivity

$$(1.2) \quad P(D)C^\infty(\Omega) = C^\infty(\Omega) ,$$

since the relation module is 0 . Thus the concept of $P(D)$ -convexity is a generalization of that of the usual $P(D)$ -convexity (see [3], [4]).

2) When $P(D) = \begin{pmatrix} D_1 \\ \vdots \\ D_n \end{pmatrix}$ is the gradient operator, the exact-

ness is just the vanishing of the first de Rham cohomology group

$$(1.3) \quad H^1(\Omega, \mathbb{C}) = 0 .$$

3) If we identify the complex space \mathbb{C}^m with \mathbb{R}^n ($n = 2m$) in the usual manner, then the pseudo-convexity of an open set Ω can be characterized by the exactness of the following sequence

$$(1.4) \quad C_{(0,0)}^{\infty}(\Omega) \xrightarrow{\bar{\partial}} C_{(0,1)}^{\infty}(\Omega) \xrightarrow{\bar{\partial}} \cdots \xrightarrow{\bar{\partial}} C_{(0,m)}^{\infty}(\Omega) \rightarrow 0$$

where $C_{(0,q)}^{\infty}$ denotes the space of all C^{∞} -forms of type $(0,q)$ defined on Ω (see [2], [5]). This can be stated as the exactness of the following sequence

$$(1.4)' \quad E_1 \xrightarrow{\bar{\partial}} E_2 \xrightarrow{\bar{\partial}} E_3$$

with $E_1 = \bigcap_{q=0}^{m-1} C_{(0,q)}^{\infty}(\Omega)$, $E_2 = \bigcap_{q=1}^m C_{(0,q)}^{\infty}(\Omega)$, $E_3 = \bigcap_{q=2}^{m+1} C_{(0,q)}^{\infty}(\Omega)$. Thus it can be written in the form (1.1).

4) When $\mathcal{P}(D)$ is quite general, we know only the following

THEOREM 1.1. (Ehrenpreis-Malgrange) *If Ω is convex, then Ω is $\mathcal{P}(D)$ -convex for any $\mathcal{P}(D)$ (See [1], [2], [5]).*

§2. Finite type systems of partial differential operators.

In the preceding section, we have explained the $\mathcal{P}(D)$ -convexity and some examples. But, other than these, no general results are known. Therefore it will have some meaning to state the following

THEOREM 2.1. *If Ω is simply connected, then Ω is $\mathcal{P}(D)$ -convex for any $\mathcal{P}(D)$ of finite type.^(*)*

Before giving a sketchy proof of the above theorem, we should explain some notions.

To a matrix $\mathcal{P}(X) = (P_{jk}(X))$ with q rows and p columns we associate the ideal $\mathcal{N} = \mathcal{N}(\mathcal{P})$ generated by all the (p,p) -minors of $\mathcal{P}(X)$. (If $p > q$, we only put $\mathcal{N} = 0$.) Let $V = V(\mathcal{P})$ be the algebraic variety defined by $\mathcal{N}(\mathcal{P})$, i.e.

$$V(\mathcal{P}) = \{z \in \mathbb{C}^n; P(z) = 0 \text{ for all } P \in \mathcal{N}(\mathcal{P})\}.$$

(*) See the definition below.

This is called the *variety attached* to the system of differential operators $\mathcal{P}(D)$.

DEFINITION. A system of partial differential operators $\mathcal{P}(D)$ is called of *finite type* if the attached variety is of dimension 0, i.e. $V(\mathcal{P})$ consists of only a finite number of points.

Now consider the following homogeneous equation

$$(2.1) \quad \mathcal{P}(D)U = 0$$

where U is an unknown element of $[C^\infty(\Omega)]^P$.

Then, using Hilbert's Nullstellensatz, it is not difficult to show the following

LEMMA The vector space over \mathbb{C} of the solutions of (2.1) is of finite dimension if and only if the system $\mathcal{P}(D)$ is of finite type. And then the solutions of (2.1) consist only of entire functions, more precisely the exponential-polynomial solutions of (2.1).

To prove Theorem 2.1, first we fix a covering of Ω by its convex open subsets: $\Omega = \bigcup_i \Omega_i$, and consider the equation

$$(2.2) \quad \mathcal{P}(D)U = F$$

where F is an arbitrary given element in $[C^\infty(\Omega)]^q$ such that

$$(2.3) \quad \mathcal{Q}(D)F = 0.$$

In each convex open set Ω_i , we can apply Theorem 1.1. and we get a solution in Ω_i . In the intersection $\Omega_i \cap \Omega_j$ of two such convex open sets, the difference of the solutions should

satisfy the homogeneous equation (2.1). Hence, according to the above lemma, we get a solution of (2.2) in the union $\Omega_i \cup \Omega_j$ by adjusting those solutions by a certain entire function. Now starting at some fixed point we can proceed along curves repeating such adjustments in a manner similar to the usual analytic continuation in function theory. The resulting solution U should be univalent according to the assumption that Ω be simply connected. This shows that (1.1) is exact. This completes the proof.

§3. Decomposition of solutions of partial differential equations.

Since partial differential operators with constant coefficients operate on $C^\infty(\Omega)$ commutatively, we see clearly that

(I) For any pair of polynomials P_1, P_2 and for any pair of functions $u_1, u_2 \in C^\infty(\Omega)$ such that

$$(3.1) \quad P_1(D)u_1 = 0, \quad P_2(D)u_2 = 0,$$

the sum

$$(3.2) \quad u = u_1 + u_2$$

satisfies the equation for the product operator

$$(3.3) \quad P_1(D)P_2(D)u = 0;$$

(II) For any polynomial P and for any multiindexed family of solutions $u_\alpha \in C^\infty(\Omega)$ ($|\alpha| \leq \nu-1$)^(*) of the equation

(*) For a multiindex $\alpha = (\alpha_1, \dots, \alpha_n)$ (a sequence of non-negative integers) we set $|\alpha| = \alpha_1 + \dots + \alpha_n$. If $x = (x_1, \dots, x_n)$ is variable point, we set $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$.

$$(3.4) \quad P(D)u_\alpha = 0$$

the sum

$$(3.5) \quad u = \sum_{|\alpha| \leq v-1} x^\alpha u_\alpha$$

is a solution of the equation for the v -times interacted operator

$$(3.6) \quad P(D)^v u = 0$$

Our question is to ask when the converses of the above facts are true, naturally assuming always that

(i) P_1 and P_2 have no common factor

for the converse of (I), and that

(ii) P is irreducible

for the converse of (II).

THEOREM 3.1. If Ω is $\mathcal{P}(D)$ -convex with $\mathcal{P}(D) = \begin{pmatrix} P_1(D) \\ P_2(D) \end{pmatrix}$, then any solution $u \in C^\infty(\Omega)$ of (3.3) can be decomposed into the form (3.2) with (3.1). Conversely, if each solution u of (3.3) can be written in the form (3.2) with (3.1) and if Ω is $P_1(D)P_2(D)$ -convex, then Ω is $\mathcal{P}(D)$ -convex.

DEFINITION. We say that an open set Ω is a *decomposing domain* if the converses of (I) and (II) are true for all P_1, P_2 and P satisfying the conditions (i) and (ii).

Application of Ehrenpreis-Malgrange's theorem thus shows

THEOREM 3.2. If Ω is convex, then Ω is a decomposing domain.

In the plane \mathbb{R}^2 , we can prove a more precise result, using Theorem 2.1.

THEOREM 3.3. An open set Ω in \mathbb{R}^2 is a decomposing domain if and only if Ω is simply connected.

The "only if" part comes from the fact that, for an open set Ω in the plane \mathbb{R}^2 , the vanishing of the first de Rham cohomology group (1.3) is equivalent to the simple-connectedness of Ω . (*)

Theorem 3.3 is no longer true for $n \geq 3$.

If we restrict ourselves to the polynomial or exponential polynomial solutions then we get, by a purely algebraic argument, the following

THEOREM 3.4. *If we assume (i) and (ii), then any polynomial (resp. exponential-polynomial) solution of (3.3) or (3.6) can be decomposed into the form (3.2) or (3.5) with u_1, u_2 or u_α polynomials (resp. exponential-polynomials).*

The content of the present article can be considered as a completion of our previous work [6]. The more detailed treatment shall be published elsewhere.

References.

- [1] Ehrenpreis, L., A fundamental principle for systems of linear differential equations with constant coefficients and some of its applications, Proc. Intern. Symp. on Linear Spaces, 161-174, Jerusalem, 1961.
- [2] Hörmander, L., An introduction to complex analysis in several variables, D. Van Nostrand, Princeton, N. J., 1966.

(*) This can be proved by a combinatorial argument, for that I thank Prof. H. TODA.

- [3] Hörmander, L., Linear partial differential operators, Springer-Verlag, 1963.
- [4] Malgrange, B., Existence et approximation des solutions des équations aux dérivées partielles et des équations de convolution, Ann. Inst. Fourier 6, 271-354. (1955).
- [5] Malgrange, B., Sur les systèmes différentiels à coefficients constants, Coll. C. N. R. S., 113-122, Paris, 1963.
- [6] Matsuura, S., Factorization of differential operators and decomposition of solutions of homogeneous equations, Osaka Math. J., 213-231, 15 (1963).